

temperature (15 to 30 °C). Add sufficient purified water to obtain 100 grams of standard sunscreen preparation.

(c) *Assay of the standard homosalate sunscreen.* Assay the standard homosalate sunscreen preparation by the following method to ensure proper concentration:

(1) *Preparation of the assay solvent.* The solvent consists of 1 percent glacial acetic acid (V/V) in denatured ethanol. The denatured ethanol should not contain a UV radiation absorbing denaturant.

(2) *Preparation of a 1-percent solution of the standard homosalate sunscreen preparation.* Accurately weigh 1 gram of the standard homosalate sunscreen preparation into a 100-milliliter volumetric flask. Add 50 milliliters of the assay solvent. Heat on a steam bath and mix well. Cool the solution to room temperature (15 to 30 °C). Then dilute the solution to volume with the assay solvent and mix well to make a 1-percent solution.

(3) *Preparation of the test solution (1:50 dilution of the 1-percent solution).* Filter a portion of the 1-percent solution through number 1 filter paper. Discard the first 10 to 15 milliliters of the filtrate. Collect the next 20 milliliters of the filtrate (second collection). Add 1 milliliter of the second collection of the filtrate to a 50-milliliter volumetric flask. Dilute this solution to volume with assay solvent and mix well. This is the test solution (1:50 dilution of the 1-percent solution).

(4) *Spectrophotometric determination.* The absorbance of the test solution is measured in a suitable double beam spectrophotometer with the assay solvent and reference beam at a wavelength near 306 nanometers.

(5) *Calculation of the concentration of homosalate.* The concentration of homosalate is determined by the following formula which takes into consideration the absorbance of the sample of the test solution, the dilution of the 1-percent solution (1:50), the weight of the sample of the standard homosalate sunscreen preparation (1 gram), and the standard absorbance value (172) of homosalate as determined by averaging the absorbance of a

large number of batches of raw homosalate:

Concentration of homosalate  

$$= \text{absorbance} \times 50 \times 100 \times 172 = \text{percent concentration by weight.}$$

#### § 352.71 Light source (solar simulator).

A solar simulator used for determining the SPF of a sunscreen drug product should be filtered so that it provides a continuous emission spectrum from 290 to 400 nanometers similar to sunlight at sea level from the sun at a zenith angle of 10° it has less than 1 percent of its total energy output contributed by nonsolar wavelengths shorter than 290 nanometers; and it has not more than 5 percent of its total energy output contributed by wavelengths longer than 400 nanometers. In addition, a solar simulator should have no significant time-related fluctuations in radiation emissions after an appropriate warmup time, and it should have good beam uniformity (within 10 percent) in the exposure plane. To ensure that the solar simulator delivers the appropriate spectrum of UV radiation, it must be measured periodically with an accurately-calibrated spectroradiometer system or equivalent instrument.

#### § 352.72 General testing procedures.

(a) *Selection of test subjects (male and female).* (1) Only fair-skin subjects with skin types I, II, and III using the following guidelines shall be selected:

*Selection of Fair-skin Subjects*  
*Skin Type and Sunburn and Tanning History*  
*(Based on first 30 to 45 minutes sun exposure after a winter season of no sun exposure.)*

I—Always burns easily; never tans (sensitive).

II—Always burns easily; tans minimally (sensitive).

III—Burns moderately; tans gradually (light brown) (normal).

IV—Burns minimally; always tans well (moderate brown) (normal).

V—Rarely burns; tans profusely (dark brown) (insensitive).

VI—Never burns; deeply pigmented (insensitive).

(2) A medical history shall be obtained from all subjects with emphasis on the effects of sunlight on their skin. Ascertain the general health of the individual, the individual's skin type (I,